CORRECTION Open Access



Correction: Eyelid and scleral thermal injury following phacoemulsification in silicone oil: a case report

Yu-Kuei Lee¹, Szu-Han Chen² and Jia-Horung Hung^{1,3*}

Correction: BMC Ophthalmol 22, 420 (2022) https://doi.org/10.1186/s12886-022-02646-x

Following publication of the original article [1], the author group has identified an error in Fig. 2. D. The correct figure is given below.

The original article [1] has been corrected.

Author details

¹Department of Ophthalmology, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan, Taiwan. ²Division of Plastic and Reconstructive Surgery, Department of Surgery, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan, Taiwan. ³Institute of Clinical Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan.

Published online: 02 December 2022

Reference

 Lee YK, Chen SH, Hung JH. Eyelid and scleral thermal injury following phacoemulsification in silicone oil: a case report. BMC Ophthalmol. 2022;22:420. https://doi.org/10.1186/s12886-022-02646-x.

The original article can be found online at https://doi.org/10.1186/s12886-022-02646-x.

*Correspondence: hungjh@mail.ncku.edu.tw

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

 $^{^3}$ Institute of Clinical Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan

Lee et al. BMC Ophthalmology (2022) 22:467 Page 2 of 2




Fig. 2 A The fragmatome tip was inserted 1 cm beneath the fluid level (arrow); **B** A high-sensitivity infrared thermal camera was used to detect the temperature of the fragmatome tip at the fluid level (arrow) of the balanced salt solution. **C** The thermal imager revealed the temperature of the fragmatome tip at the fluid level (arrow) of the silicone oil. **D** The change in the temperature at the fragmatome tip in the balanced salt solution and silicone oil in 240 s