# CORRECTION Open Access



# Correction: Efficacy of topical 0.05% cyclosporine A and 0.1% sodium hyaluronate in post-refractive surgery chronic dry eye patients with ocular pain

Lu Zhao<sup>1†</sup>, Jiawei Chen<sup>2†</sup>, Hongyu Duan<sup>1</sup>, Tingting Yang<sup>3</sup>, Baikai Ma<sup>1</sup>, Yifan Zhou<sup>1</sup>, LinBo Bian<sup>1</sup>, Xiying Cai<sup>4</sup> and Hong Qi<sup>1,2\*</sup>

Correction: BMC Ophthalmol 24, 28 (2024) https://doi.org/10.1186/s12886-024-03294-z

Following publication of the original article [1], the authors would like to replace Figure 3 as it mistakenly uses the bar graph from Figure 2.

Figure 2 remains accurate and unaffected.

This error does not impact the conclusions of the article.

The correct and incorrect version of Fig. 3 can be found below:

<sup>†</sup>Lu Zhao and Jiawei Chen contributed equally to this work.

The online version of the original article can be found at https://doi.org/10.1186/s12886-024-03294-z.

\*Correspondence:

Hong Qi

doctorqihong@163.com

<sup>1</sup>Department of Ophthalmology, Beijing Key Laboratory of Restoration of Damaged Ocular Nerve, Peking University Third Hospital, Beijing, China <sup>2</sup>Institute of Medical Technology, Peking University Health Science Center, 49 North Garden Road, Haidian District, Beijing 100191, China

<sup>3</sup>Department of Ophthalmology, Guangdong Provincial People's Hospital (Guangdong Academy of Medical Sciences, Southern Medical University, Guangzhou, China

<sup>4</sup>Peking University First Hospital, Beijing, China



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>. The Creative Commons Public Domain Dedication waiver (<a href="http://creativecommons.org/publicdomain/zero/1.0/">http://creativecommons.org/publicdomain/zero/1.0/</a>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Zhao et al. BMC Ophthalmology (2024) 24:223 Page 2 of 3

### Correct




Fig. 3 The proportion of ocular pain severity in post–refractive surgery DED patients with ocular pain before and after treatment. The severity of ocular pain was scored according to Numerical Rating Scale (NRS) (range, 0–10)

### Incorrect




Fig. 3 The proportion of ocular pain severity in post–refractive surgery DED patients with ocular pain before and after treatment. The severity of ocular pain was scored according to Numerical Rating Scale (NRS) (range, 0–10)

The original article has been updated.

Zhao et al. BMC Ophthalmology (2024) 24:223 Page 3 of 3

Published online: 28 May 2024

## **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

# References

 Zhao L, Chen J, Duan H, et al. Efficacy of topical 0.05% cyclosporine A and 0.1% sodium hyaluronate in post-refractive surgery chronic dry eye patients with ocular pain. BMC Ophthalmol. 2024;24:28. https://doi.org/10.1186/ s12886-024-03294-z.