CORRECTION Open Access

Correction to: Thickness profiles of the corneal epithelium along the steep and flat meridians of astigmatic corneas after orthokeratology



Jiaqi Zhou^{1†}, Feng Xue^{1†}, Xingtao Zhou¹, Rajeev Krishnan Naidu² and Yishan Qian^{1*}

Correction to: BMC Ophthalmol 20, 240 (2020) https://doi.org/10.1186/s12886-020-01477-y

Following publication of the original article [1], we were notified that the authors would like to remove affiliation 2 (Sydney Eye Hospital, Sydney, NSW, 2000, Australia) from the author Xingtao Zhou and Yishan Qian, and affiliation 1 (Department of Ophthalmology, Eye and ENT Hospital, Fudan University; Key Laboratory of Myopia of The State Health Ministry, 200031, 83 Fenyang Road, Shanghai, People's Republic of China) from the author Rajeev Krishnan Naidu.

Author details

¹Department of Ophthalmology, Eye and ENT Hospital, Fudan University; Key Laboratory of Myopia of The State Health Ministry, 200031, 83 Fenyang Road, Shanghai, People's Republic of China. ²Sydney Eye Hospital, Sydney, NSW 2000. Australia

Published online: 18 August 2020

Reference

 Zhou, et al. Thickness profiles of the corneal epithelium along the steep and flat meridians of astigmatic corneas after orthokeratology. BMC Ophthalmol. 2020;20:240. https://doi.org/10.1186/s12886-020-01477-y.

The original article can be found online at https://doi.org/10.1186/s12886-020-01477-y.

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*} Correspondence: thronebird31@hotmail.com

[†]Jiaqi Zhou and Feng Xue These authors contributed equally to this manuscript and should be considered co-first authors.

¹Department of Ophthalmology, Eye and ENT Hospital, Fudan University; Key Laboratory of Myopia of The State Health Ministry, 200031, 83 Fenyang Road, Shanghai, People's Republic of China